arXiv:1109.3618v1 [math.AP] 16 Sep 2011

Existence and asymptotic behaviour of solutions of the very fast diffusion equation

Shu-Yu Hsu
Department of Mathematics
National Chung Cheng University
168 University Road, Min-Hsiung
Chia-Yi 621, Taiwan, R.O.C.
e-mail: syhsu@math.ccu.edu.tw

Sept 16, 2011

Abstract

Let $n\ge 3$, $0<m\le(n-2)/n$, $p>\max(1,(1-m)n/2)$, and $0\le u_0\in L^p_{loc}(\mathbb{R}^n)$ satisfy $\liminf_{R\to\infty}R^{-n+\frac{2}{1-m}}\int_{|x|\le R}u_0\,dx=\infty$. We prove the existence of unique global classical solution of $u_t=\frac{n-1}{m}\Delta u^m$, $u>0$, in $\mathbb{R}^n\times(0,\infty)$, $u(x,0)=u_0(x)$ in $\mathbb{R}^n$. If in addition $0<m<(n-2)/n$ and $u_0(x)\approx A|x|^{-q}$ as $|x|\to\infty$ for some constants $A>0$, $q<n/p$, we prove that there exist constants α, β, such that the function $v(x,t)=t^{\alpha}u(t^{\beta}x,t)$ converges uniformly on every compact subset of $\mathbb{R}^n$ to the self-similar solution $\psi(x,1)$ of the equation with $\psi(x,0)=A|x|^{-q}$ as $t\to\infty$. Note that when $m=(n-2)/(n+2)$, $n\ge 3$, if $g_{ij}=u^{\frac{4}{n+2}}\delta_{ij}$ is a metric on $\mathbb{R}^n$ that evolves by the Yamabe flow $\partial g_{ij}/\partial t=-Rg_{ij}$ with $u(x,0)=u_0(x)$ in $\mathbb{R}^n$ where R is the scalar curvature, then $u(x,t)$ is a global solution of the above fast diffusion equation.

Key words: existence, global solution, very fast diffusion equation, asymptotic behaviour, Yamabe flow

AMS Mathematics Subject Classification: Primary 35K15, 35B40 Secondary 35K65, 58J35

1 Introduction

Recently there is a lot of study of the equation [A], [BBDGV], [DK], [DP], [DS1], [DS2], [Hs2], [P], [V1–3],

$$u_t=\frac{n-1}{m}\Delta u^m. \tag{1.1}$$

The above equation arises in many physical models and in geometry. When $m > 1$, (1.1) models the flow of gases through porous media [A]. When $m = 1$, (1.1) is the heat equation after a rescaling. When $m = -1$, the equation (1.1) arises in the model of heat conduction in solid hydrogen [R]. As observed by M. Del Pino, M. Sáez, P. Daskalopoulos and N. Sesum [PS], [DS2], when $m = (n-2)/(n+2)$, $n \geq 3$, if $g_{ij} = u^{\frac{4}{n+2}}\delta_{ij}$ is a metric on $\mathbb{R}^n$ that evolves by the Yamabe flow

$$\frac{\partial}{\partial t} g_{ij} = -R g_{ij} \tag{1.2}$$

where R is the scalar curvature, then u satisfies (1.1). We refer the readers to the book [V3] by J.L. Vazquez for an introduction of the equation (1.1) and the books [DK] by P. Daskalopoulos and C.E. Kenig, [V2] by J.L. Vazquez, for the most recent results on (1.1).

As observed by L.A. Peletier in [P] (Theorem 11.3 of [P]) the behaviour of the solutions of (1.1) for $(n-2)/n < m < 1$, $n \geq 3$, are very different from the behaviour of the solutions of (1.1) for $0 < m < (n-2)/n$, $n \geq 3$. For any $0 < m < 1$, $n \geq 3$, let $0 \leq u_0 \in L^1(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$ for some constant p satisfying

$$p > \max(1, (1-m)n/2). \tag{1.3}$$

By Theorem 11.3 of [P] for any $0 < m < (n-2)/n$ and $n \geq 3$ there exists a constant $T > 0$ and a distribution solution $u \in C([0,T]; L^1(\mathbb{R}^n)) \cap L^\infty([\delta, T] \times \mathbb{R}^n)$ for any $\delta \in (0, T]$ of (1.1) with $u(x,0) = u_0(x)$ on $\mathbb{R}^n$ which vanishes identically at time T. For example for any $k > 0$ and $T > 0$, the Barenblatt solution ([DS2]),

$$B_k(x,t) = \left(\frac{C_*}{k + (T-t)_+^{\frac{2}{n-2-nm}}|x|^2}\right)^{\frac{1}{1-m}} (T-t)_+^{\frac{n}{n-2-nm}}, \tag{1.4}$$

where $C_* = 2(n-1)(n-2-nm)/(1-m)$ is a classical solution of (1.1) in $\mathbb{R}^n \times (0,T)$ which vanishes identically at time T. On the other hand an examination of the proof of [HP] shows that the existence proof of global solutions of

$$\begin{cases} u_t = \dfrac{n-1}{m}\Delta u^m, u > 0, & \text{in } \mathbb{R}^n \times (0,T) \\ u(x,0) = u_0(x) & \text{in } \mathbb{R}^n \end{cases} \tag{1.5}$$

is valid only for $(n-2)/n < m < 1$ and $n \geq 3$. Then by the result of [HP] for $n \geq 3$, $(n-2)/n < m < 1$, and any $0 \leq u_0 \in L^1_{loc}(\mathbb{R}^n)$, $u_0 \not\equiv 0$, there exists a unique global solution of (1.5) in $\mathbb{R}^n \times (0,\infty)$.

In this paper we will prove that if $n \geq 3$, $0 < m \leq (n-2)/n$, and $0 \leq u_0 \in L^p_{loc}(\mathbb{R}^n)$ for some constant p satisfying (1.3) and

$$\liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\leq R} u_0\, dx = \infty, \tag{1.6}$$

then (1.5) has a global solution u in $\mathbb{R}^n \times (0,\infty)$. When u_0 is radially symmetric, the condition (1.6) is also shown in [DP] to imply global existence of solution of (1.5) for

$m < 0$. On the other hand when $m > 1$, by the result of Aronson and Caffarelli [AC], if (1.5) has a global solution in $\mathbb{R}^n \times (0, \infty)$, then

$$\limsup_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le R} u_0\, dx = 0.$$

Thus our result extends the result of [AC] and [DP]. Let

$$\alpha = \frac{q}{2 - q(1-m)} \quad \text{and} \quad \beta = \frac{1}{2 - q(1-m)}. \tag{1.7}$$

We will also prove that when $u_0(x) \approx A|x|^{-q}$ for some constants $A > 0$, $q < n/p$, as $|x| \to \infty$ where p satisfies (1.3), then under some mild condition on u_0 the rescaled function

$$v(x,t) = t^{\alpha} u(t^{\beta} x, t) \tag{1.8}$$

converges uniformly on every compact subset of $\mathbb{R}^n$ to the self-similar solution $\psi(x,1)$ of (1.1) with $\psi(x,0) = A|x|^{-q}$ as $t \to \infty$. The function $\widetilde{v}(x) = \psi(x,1)$ is radially symmetric and satisfies the elliptic equation

$$\frac{n-1}{m}\Delta\widetilde{v}^m + \alpha\widetilde{v} + \beta x \cdot \nabla\widetilde{v} = 0, \widetilde{v} > 0, \quad \text{in } \mathbb{R}^n. \tag{1.9}$$

The main results we obtain in this paper are the following.

Theorem 1.1. *Let $n \ge 3$, $0 < m \le (n-2)/n$, and let p satisfy* (1.3)*. Then there exists a constant $C_1 > 0$ such that if $0 \le u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies*

$$\liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le R} u_0\, dx \ge C_1 T^{\frac{1}{1-m}} \tag{1.10}$$

for some constant $T > 0$, then there exists a unique positive solution $u \in C^{\infty}(\mathbb{R}^n \times (0,\infty))$ of (1.5) *in $\mathbb{R}^n \times (0,T)$ which satisfy*

$$u_t \le \frac{u}{(1-m)t} \tag{1.11}$$

in $\mathbb{R}^n \times (0,T)$.

Theorem 1.2. *Let $n \ge 3$, $0 < m \le (n-2)/n$, and let p satisfy* (1.3)*. Suppose $0 \le u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies* (1.6)*. Then* (1.5) *has a unique global positive solution $u \in C^{\infty}(\mathbb{R}^n \times (0,\infty))$ which satisfies* (1.11) *in $\mathbb{R}^n \times (0,\infty)$.*

Theorem 1.3. *Let $n \ge 3$, $0 < m < (n-2)/n$, and $q < n/p$ for some constant p satisfying* (1.3)*. Suppose $0 \le u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies $u_0 = \overline{u}_0 + \phi$ where $0 \le \overline{u}_0 \in L^p_{loc}(\mathbb{R}^n)$ and $\phi \in L^1(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$ such that*

$$\lim_{|x|\to\infty} |x|^q \overline{u}_0(x) = A \tag{1.12}$$

for some constant $A > 0$. Let α and β be given by (1.7)*. Let u be the unique global solution of* (1.5) *in $\mathbb{R}^n \times (0,\infty)$ given by Corollary 2.8 which satisfies* (1.11) *in $\mathbb{R}^n \times (0,\infty)$. Let v be given by* (1.8)*. Then the rescaled function v converges uniformly on every compact subset of $\mathbb{R}^n$ to the unique radially symmetric self-similar solution $\psi(x,1)$ of* (1.1) *with $\psi(x,0) = A|x|^{-q}$ as $t \to \infty$.*

Corollary 1.4. *Let* $n \ge 3$ *and* $m = (n-2)/(n+2)$. *Let* p, q, α, β, u_0, v, *be as in Theorem 1.3. Suppose the metric* $g_{ij} = u^{4/(n+2)}\delta_{ij}$ *evolves by the Yamabe flow* (1.2) *with* $u(x,0) = u_0(x)$ *on* $\mathbb{R}^n$. *Then* $v(x,t)$ *converges uniformly on every compact subset of* $\mathbb{R}^n$ *to the unique radially symmetric self-similar solution* $\psi(x,1)$ *of* (1.1) *with* $\psi(x,0) = A|x|^{-q}$ *as* $t \to \infty$.

Note that the Barenblatt solution $B_k(x,t)$ given by (1.4) satisfies

$$\lim_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le R} B_k(x,0)\,dx = \frac{\omega_n C_*^{\frac{1}{1-m}} T^{\frac{1}{1-m}}}{n-2(1-m)^{-1}}$$

where ω_n is the surface area of the unit sphere S^{n-1} in $\mathbb{R}^n$ and it vanishes in a finite time T. Hence Theorem 1.1 is sharp.

The plan of the paper is as follows. In section two we will prove the existence of global solutions of (1.5). In section three we will prove the asymptotic large time behaviour of the global solution of (1.5).

We first start will some definitions (cf. [Hu2]). For any $x_0 \in \mathbb{R}^n$, $R > 0$, $T > 0$, let $B_R(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < R\}$, $B_R = B_R(0)$, $Q_R(x_0) = B_R(x_0) \times (0,\infty)$, $Q_R = Q_R(0)$, $Q_R^T(x_0) = B_R(x_0) \times (0,T)$ and $Q_R^T = Q_R^T(0)$. For any set $A \subset \mathbb{R}^n$, we let χ_A be the characteristic function of the set A. For any domain $\Omega \subset \mathbb{R}^n$, $T > 0$, $0 < m < 1$, we say that u is a solution (subsolution, supersolution) of (1.1) in $\Omega \times (0,T)$ if $u > 0$ in $\Omega \times (0,T)$ and is a classical solution (subsolution, supersolution) of (1.1) in $\Omega \times (0,T)$. For any $0 \le u_0 \in L^1_{loc}(\Omega)$ we say that a solution u of (1.1) in $\Omega \times (0,T)$ has initial value u_0 if $\|u(\cdot,t) - u_0\|_{L^1(K)} \to 0$ as $t \to 0$ for any compact subset $K \subset \Omega$.

For any bounded smooth domain $\Omega \subset \mathbb{R}^n$, $0 \le u_0 \in L^1_{loc}(\Omega)$, and $0 \le g \in L^1(\partial\Omega \times (0,T))$, we say that u is a solution of the Dirichlet problem

$$\begin{cases} u_t = \dfrac{n-1}{m}\Delta u^m & \text{in } \Omega \times (0,T) \\ u = g & \text{on } \partial\Omega \times (0,T) \\ u(x,0) = u_0(x) & \text{in } \Omega \end{cases} \tag{1.13}$$

if u is a positive classical solution of (1.1) in $\Omega \times (0,T)$ with initial value u_0 and satisfies

$$\int_{t_1}^{t_2}\int_\Omega \left(\frac{n-1}{m}u^m\Delta\eta + u\eta_t\right)dx\,ds = \frac{n-1}{m}\int_{t_1}^{t_2}\int_{\partial\Omega} g^m \frac{\partial\eta}{\partial n}\,d\sigma dt + \int_\Omega u(x,t_2)\,dx - \int_\Omega u(x,t_1)\,dx \tag{1.14}$$

for any $0 < t_1 < t_2 < T$ and $\eta \in C^2(\overline{\Omega} \times (0,T))$ satisfying $\eta = 0$ on $\partial\Omega \times (0,T)$ where $\partial/\partial n$ is the exterior normal derivative on $\partial\Omega$. We say that u is a weak solution of the Dirichlet problem (1.13) if $0 \le u \in C([0,T_R); L^1(\Omega))$ satisfies (1.14) for any $0 < t_1 < t_2 < T$ and $\eta \in C^2(\overline{\Omega} \times (0,T))$ satisfying $\eta = 0$ on $\partial\Omega \times (0,T)$ and u has initial value u_0.

We say that u is a solution of the Dirichlet problem

$$\begin{cases} u_t = \dfrac{n-1}{m}\Delta u^m, u > 0, & \text{in } \Omega \times (0,\infty) \\ u(x,t) = \infty & \text{on } \partial\Omega \times (0,\infty) \\ u(x,0) = u_0(x) & \text{in } \Omega \end{cases} \tag{1.15}$$

if u is a positive classical solution of (1.1) in $\Omega \times (0, \infty)$ with initial value u_0 and

$$\lim_{\substack{(y,s)\to(x,t)\\(y,s)\in\Omega\times(0,\infty)}} u(y,s) = \infty \quad \forall (x,t) \in \partial\Omega \times (0,\infty)$$

We say that u is a solution (subsolution, supersolution) of (1.1) in $\mathbb{R}^n \times (0,T)$ if $u > 0$ in $\mathbb{R}^n \times (0,T)$ and is a classical solution (subsolution, supersolution) of (1.1) in $\mathbb{R}^n \times (0,T)$. For any $0 \le u_0 \in L^1_{loc}(\mathbb{R}^n)$ we say that a solution u of (1.1) in $\mathbb{R}^n \times (0,T)$ has initial value u_0 if $\|u(\cdot,t) - u_0\|_{L^1(K)} \to 0$ as $t \to 0$ for any compact subset $K \subset \mathbb{R}^n$.

We will assume that $n \ge 3$, $0 < m \le (n-2)/n$, and (1.3) hold for the rest of the paper.

2 Existence of global solutions

In this section we will prove the existence of global solutions of (1.5). We first extend some results of [Hu2]. We first observe that by (1.3),

$$\frac{2}{n} + \frac{\alpha+m}{\alpha+1} \ge \frac{2}{n} + \frac{p-1+m}{p} > 1 \quad \forall \alpha \ge p-1.$$

Hence there exists a constant

$$1 < k' < \frac{1}{q} + \frac{\alpha+m}{\alpha+1} \quad \forall \alpha \ge p-1 \tag{2.1}$$

where $q = n/2$. Let

$$k = \left(\frac{\alpha+m}{\alpha+1}\right)k'. \tag{2.2}$$

Then by the same argument as the proof of Theorem 1.6 of [Hu2] but with the α_0, k, k', there being replaced by $\alpha_0 = p-1$ and k, k', given by (2.1), (2.2), the proof of Theorem 1.6 of [Hu2] remains valid for $n \ge 3$, $0 < m \le (n-2)/n$, and p satisfying (1.3). Hence we have the following theorem.

Theorem 2.1. *(cf. Theorem 1.6 of [Hu2]) Let $n \ge 3$, $0 < m \le (n-2)/n$, and let p satisfy* (1.3). *Suppose u is a solution of* (1.1) *in $Q^* = \{(x,t) \in \mathbb{R}^{n+1} : |x| < 2, -4 < t < 0\}$. Let $Q = \{(x,t) \in \mathbb{R}^{n+1} : |x| < 1, -1 < t < 0\}$. Then there exist constants $C > 0$ and $\theta > 0$ such that*

$$\|u\|_{L^\infty(Q)} \le C\left(1 + \iint_{Q^*} u^p \, dx \, dt\right)^{\frac{\theta}{p}}.$$

We next observe that the result of Lemma 1.7 and Lemma 1.9 of [Hu2] remains valid for any $n \ge 3$, $0 < m < 1$, p satisfying (1.3), and $0 \le g \in L^\infty(\partial\Omega \times (0,\infty))$ where Ω is a bounded smooth domain. By Lemma 1.7 and Lemma 1.9 of [Hu2], Theorem 2.1, and an argument similar to that of [Hu2] we have the following three results.

Corollary 2.2. *(cf. Corollary 1.8 of [Hu2]) Let $n \geq 3$, $0 < m \leq (n-2)/n$, and let p satisfy* (1.3). *Suppose u is a solution of* (1.1) *in $\Omega \times (0,T)$ with initial value $0 \leq u_0 \in L^p_{loc}(\Omega)$. Then for any $B_{R_1}(x_0) \subset \overline{B_{R_2}(x_0)} \subset \Omega$ and $0 < t_1 < T$ there exist constants $C > 0$ and $\theta > 0$ such that*

$$\|u\|_{L^\infty(B_{R_1}(x_0)\times[t_1,T))} \leq C\left(1 + \int_{B_{R_2}(x_0)} u_0^p\,dx\right)^{\frac{\theta}{p}}.$$

Theorem 2.3. *(cf. Corollary 1.11 of [Hu2]) Let $n \geq 3$, $0 < m \leq (n-2)/n$, and let p satisfy* (1.3). *Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain and let $0 \leq g \in L^\infty(\partial\Omega \times (0,T))$. Suppose u is a solution of* (1.13) *in $\Omega \times (0,T)$. Then for any $0 < t_1 < T$ there exist constants $C > 0$ and $\theta > 0$ such that*

$$\|u\|_{L^\infty(\overline{\Omega}\times[t_1,T))} \leq C\left(k_g^p|\Omega| + \int_\Omega u_0^p\,dx\right)^{\frac{\theta}{p}} + k_g$$

where $k_g = \max(1, \|g\|_{L^\infty(\partial\Omega\times(0,T))})$.

Theorem 2.4. *(cf. Theorem 2.5 and Theorem 2.11 of [Hu2]) Let $n \geq 3$, $0 < m \leq (n-2)/n$, and let p satisfy* (1.3). *Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain. Then there exists a minimal solution u for* (1.15) *which satisfies* (1.11) *in $\Omega \subset \mathbb{R}^n$. When Ω is a smooth bounded star shape domain, the solution u for* (1.15) *is unique.*

Note that a result similar to Corollary 2.2 is also obtained recently by M. Bonforte and J.L. Vazquez (Theorem 2.1 of [BV]).

Lemma 2.5. *Let $n \geq 3$, $0 < m \leq (n-2)/n$, and let p satisfy* (1.3). *Suppose $0 \leq u_0 \in L^p_{loc}(\mathbb{R}^n)$. For any $R > 0$ let $0 \leq w_R \in C([0,T_R); L^1(B_{5R}))$ be the unique weak solution of*

$$\begin{cases} w_t = \dfrac{n-1}{m}\Delta w^m & \text{in } Q_{5R}^{T_R} \\ w = 0 & \text{on } \partial B_{5R} \times (0,T_R) \\ w(x,0) = u_0(x)\chi_{B_{2R}}(x) & \text{in } B_{5R} \end{cases} \tag{2.3}$$

given by [BC] (cf. P.537 of [BV]) which extincts in a finite time $T_R > 0$. Then there exists a constant $C_1 > 0$ such that if (1.10) *holds for some constant $T > 0$, then there exist constants $0 < \delta < 1$, $R_0 > 0$, such that for any $R \geq R_0$, $T_R > T$, w_R is continuous on $\overline{B_{\delta R}} \times (0,T]$ and*

$$\inf_{\overline{B_{\delta R}}\times[t_1,T]} w_R > 0 \quad \forall 0 < t_1 < T, R \geq R_0. \tag{2.4}$$

Proof: Suppose there exists $T > 0$ such that u_0 satisfies (1.10) for some constant $C_1 > 0$ to be determined later. For any $\varepsilon > 0$ let $w_{R,\varepsilon}$ be the solution of

$$\begin{cases} w_t = \dfrac{n-1}{m}\Delta w^m & \text{in } Q_{5R} \\ w = \varepsilon & \text{on } \partial B_{5R} \times (0,\infty) \\ w(x,0) = u_0(x)\chi_{B_{2R}}(x) + \varepsilon & \text{in } B_{5R} \end{cases}$$

Then $w_{R,\varepsilon} \ge w_{R,\varepsilon'} \ge \varepsilon'$ in Q_{5R} for any $\varepsilon > \varepsilon' > 0$ and $w_{R,\varepsilon}$ decreases to the weak solution w_R of (2.3) as $\varepsilon \to 0$. By (1.10) there exists a constant $0 < \delta < 1$ such that

$$(1-\delta)^{n-\frac{2}{1-m}} \liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le R} u_0\,dx > (3C_1/4)T^{\frac{1}{1-m}}$$

Let $|x_0| \le \delta R$. Then

$$\begin{aligned}\liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x-x_0|\le R} u_0\,dx \ge& \liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le(1-\delta)R} u_0\,dx\\ \ge&(1-\delta)^{n-\frac{2}{1-m}} \liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{|x|\le R} u_0\,dx > (3C_1/4)T^{\frac{1}{1-m}}.\end{aligned} \tag{2.5}$$

Let $0 \le v_R \in C([0,\widetilde{T}_R); L^1(B_{3R}))$ be the unique weak solution of

$$\begin{cases} v_t = \dfrac{n-1}{m}\Delta v^m & \text{in } Q_{3R}^{\widetilde{T}_R}(x_0)\\ v = 0 & \text{on } \partial B_{3R}(x_0)\times(0,\widetilde{T}_R)\\ v(x,0) = u_0(x)\chi_{B_R(x_0)}(x) & \text{in } B_{3R}(x_0)\end{cases} \tag{2.6}$$

given by [BC] (cf. P.537 of [BV]) which extincts in a finite time $\widetilde{T}_R > 0$ and let $v_{R,\varepsilon}$ be the solution of

$$\begin{cases} v_t = \dfrac{n-1}{m}\Delta v^m & \text{in } Q_{3R}(x_0)\\ v = \varepsilon & \text{on } \partial B_{3R}(x_0)\times(0,\infty)\\ v(x,0) = u_0(x)\chi_{B_R(x_0)}(x) + \varepsilon & \text{in } B_{3R}(x_0)\end{cases}$$

for any $\varepsilon > 0$. Since $v_{R',\varepsilon} \ge \varepsilon$ in $Q_{3R'}(x_0)$ for any $R' > 0$, by the maximum principle (cf. Lemma 2.3 of [DaK]),

$$v_{R',\varepsilon} \ge v_{R,\varepsilon} \quad \text{in } Q_{3R}(x_0) \quad \text{for any } R' > R > 0 \tag{2.7}$$

and

$$w_{R,\varepsilon} \ge v_{R,\varepsilon} \quad \text{in } Q_{3R}(x_0) \quad \forall R > 0, \varepsilon > 0. \tag{2.8}$$

Since $v_{R,\varepsilon}$ decreases to v_R as $\varepsilon \to 0$, by (2.8),

$$w_{R,\varepsilon} \ge v_R \quad \text{in } Q_{3R}^{\widetilde{T}_R}(x_0). \tag{2.9}$$

Letting $\varepsilon \to 0$ in (2.7) we get $v_{R'} \ge v_R$ in $Q_{3R}^{\widetilde{T}_R}(x_0)$ for any $R' > R > 0$. Hence $\widetilde{T}_{R'} \ge \widetilde{T}_R$ for any $R' > R > 0$. By (1.18) of [BV] there exists a constant $C_2 > 0$ such that

$$\widetilde{T}_R \ge C_2R^2\left[\frac{\int_{B_R(x_0)} u_0\,dx}{Vol(B_{3R}\setminus B_{2R})}\right]^{1-m} \qquad \forall R > 0. \tag{2.10}$$

Letting $R \to \infty$ in (2.10), by (2.5) we get

$$\lim_{R\to\infty} \widetilde{T}_R^{\frac{1}{1-m}} \ge \frac{3nC_1C_2^{\frac{1}{1-m}}}{4(3^n - 2^n)\omega_n} T^{\frac{1}{1-m}}$$

where ω_n is the surface area of the unit sphere S^{n-1} in $\mathbb{R}^n$. We now choose $C_1 \ge 4(3^n - 2^n)\omega_n/(nC_2^{1/(1-m)})$. Then

$$\lim_{R\to\infty} \widetilde{T}_R > T. \tag{2.11}$$

By (1.28) of [BV] there exist constants $C_3 > 0$, $C_4 > 0$, such that

$$\frac{1}{R^{n-\frac{2}{1-m}}} \int_{B_R(x_0)} u_0 \, dx \le C_3 t^{\frac{1}{1-m}} + C_4 \widetilde{T}_R^{\frac{1}{1-m}} R^{\frac{2m}{1-m}} t^{-\frac{m}{1-m}} v_R^m(t, x_0) \quad \forall 0 < t < \widetilde{T}_R, R > 0. \tag{2.12}$$

Let $C_1 = \max(3C_3, 4(3^n - 2^n)\omega_n/(nC_2^{1/(1-m)}))$. By (2.5),

$$\liminf_{R\to\infty} \frac{1}{R^{n-\frac{2}{1-m}}} \int_{B_R(x_0)} u_0 \, dx \ge (9C_3/4)T^{\frac{1}{1-m}}. \tag{2.13}$$

Then by (2.11) and (2.13) there exists a constant $R_0 > 0$ such that $\widetilde{T}_R > T$ for any $R \ge R_0$ and

$$\frac{1}{R^{n-\frac{2}{1-m}}} \int_{B_R(x_0)} u_0 \, dx > 2C_3 T^{\frac{1}{1-m}} \quad \forall R \ge R_0. \tag{2.14}$$

By (2.9), (2.12), and (2.14),

$$C_4 \widetilde{T}_R^{\frac{1}{1-m}} R^{\frac{2m}{1-m}} t^{-\frac{m}{1-m}} w_{R,\varepsilon}^m(x_0, t) \ge C_4 \widetilde{T}_R^{\frac{1}{1-m}} R^{\frac{2m}{1-m}} t_0^{-\frac{m}{1-m}} v_R^m(x_0, t) > C_3(2T^{\frac{1}{1-m}} - t^{\frac{1}{1-m}}) > 0$$

for any $R \ge R_0$, $\varepsilon > 0$, $|x_0| \le \delta R$ and $0 < t \le T$. Hence

$$\inf_{\overline{B_{\delta R}}\times[t_1,T]} w_{R,\varepsilon}^m(x, t) \ge (C_3/C_4)\widetilde{T}_R^{-\frac{1}{1-m}} R^{-\frac{2m}{1-m}} t_1^{\frac{m}{1-m}} (2T^{\frac{1}{1-m}} - t_1^{\frac{1}{1-m}}) > 0 \quad \forall 0 < t_1 < T, R \ge R_0, \varepsilon > 0. \tag{2.15}$$

By Corollary 2.2 for any $R \ge R_0$, $T \ge t_1 > 0$, there exist constants $C_R > 0$ and $\theta > 0$ such that

$$\|w_{R,\varepsilon}\|_{L^\infty(\overline{B_{\delta R}}\times[t_1,T])} \le C_R \left(1 + \int_{B_{2R}} u_0^p \, dx\right)^{\frac{\theta}{p}} \quad \forall 0 < \varepsilon < 1. \tag{2.16}$$

Let $R \ge R_0$. By (2.15) and (2.16) the equation (1.1) for the sequence $\{w_{R,\varepsilon}\}_{0<\varepsilon<1}$ is uniformly parabolic on $\overline{B_{\delta R}} \times [t_1, T]$. By reducing δ if necessary by the parabolic Schauder estimates [LSU] the sequence $\{w_{R,\varepsilon}\}_{0<\varepsilon<1}$ is equi-Holder continuous in $C^2(K)$ on every compact subset $K \subset \overline{B_{\delta R}} \times [t_1, T]$. Hence by the Ascoli Theorem $w_{R,\varepsilon}$ converges in C^2 on every compact subset of $\overline{B_{\delta R}} \times (0, T]$ to w_R as $\varepsilon \to 0$. Hence $w_R \in C^2(\overline{B_{\delta R}} \times (0, T])$. Letting $\varepsilon \to 0$ in (2.15), we get (2.4) and the lemma follows. □

Proof of Theorem 1.1: Uniqueness of solution of (1.5) is given by Theorem 2.3 of [HP]. Hence we will only need to prove existence of solution of (1.5). We will give two different methods of construction of solution of (1.5). Let $C_1 > 0$ be as in Lemma 2.5.

First method: For any $R > 0$ by Theorem 2.4 (1.15) has a unique solution u_R in Q_R with $\Omega = B_R$ which satisfies (1.11) in Q_R. By the maximal principle (Lemma 2.9 of [Hu2]),

$$u_R \ge u_{R'} > 0 \quad \text{in } Q_R \quad \forall R' > R > 0. \tag{2.17}$$

Hence

$$0 \le u(x,t) = \lim_{R\to 0} u_R(x,t) \quad \forall (x,t) \in \mathbb{R}^n \times (0,\infty)$$

exists. By Corollary 2.2 for any $R_1 > 0$, $t_2 > t_1 > 0$, there exist constants $C > 0$ and $\theta > 0$ such that

$$\|u_R\|_{L^\infty(\overline{B_{R_1}(x_0)}\times[t_1,t_2])} \le C_{R_1,t_1,t_2} := C\left(1 + \int_{B_{2R_1}(x_0)} u_0^p\,dx\right)^{\frac{\theta}{p}} \quad \forall R > 3R_1 + 1 \tag{2.18}$$

$$\Rightarrow \quad \|u\|_{L^\infty(\overline{B_{R_1}(x_0)}\times[t_1,t_2])} \le C_{R_1,t_1,t_2} \qquad \text{as } R \to \infty. \tag{2.19}$$

We claim that

$$\inf_{\overline{B_R}\times[t_1,T]} u(x,t) > 0 \quad \forall R > 0, T > t_1 > 0. \tag{2.20}$$

Suppose the claim is false. Let R_0, w_R, T_R, be as in Lemma 2.5. For any $\varepsilon > 0$ let $w_{R,\varepsilon}$ be as in the proof of Lemma 2.5. By the maximum principle (Lemma 2.3 of [DaK] and Lemma 2.9 of [Hu2]),

$$\begin{aligned} & w_{R_1',\varepsilon} \ge w_{R_1,\varepsilon} \quad \text{in } Q_{5R_1}^{T_{R_1}} \quad \forall R_1' > R_1 > 0, \varepsilon > 0 \\ \Rightarrow \quad & w_{R_1'} \ge w_{R_1} \quad \text{in } Q_{5R_1}^{T_{R_1}} \quad \forall R_1' > R_1 > 0 \quad \text{as } \varepsilon \to 0 \end{aligned} \tag{2.21}$$

and

$$u_R \ge w_{R_1} \quad \text{in } Q_{5R_1}^{T_{R_1}} \quad \forall R > 5R_1 > 0. \tag{2.22}$$

Letting $R \to \infty$ in (2.22),

$$u \ge w_{R_1} \quad \text{in } Q_{5R_1}^{T_{R_1}} \quad \forall R_1 > 0. \tag{2.23}$$

By (2.4) and (2.23) the claim (2.20) holds. By (2.18), (2.20), the equation (1.1) for the sequence $\{u_R\}_{R>R_0}$ are uniformly parabolic on every compact subset of $\mathbb{R}^n \times (0,T]$. By the Schauder estimates for parabolic equations [LSU] the sequence $\{u_R\}_{R>R_0}$ are equi-Holder continuous on every compact subset of $\mathbb{R}^n \times (0,T]$. Hence by the Ascoli Theorem and (2.17) the sequence u_R decreases and converges uniformly on every compact subset of $\mathbb{R}^n \times (0,T]$ to a solution u of (1.1) in $\mathbb{R}^n \times (0,T)$ as $R \to \infty$. Then u also satisfies (1.11) in $\mathbb{R}^n \times (0,T)$.

We will now prove that u has initial value u_0. By the Kato inequality [K] (cf. [DS1]) and an argument similar to the proof of Lemma 3.1 of [HP] and Lemma 2.3 of [Hu2],

$$\begin{aligned}\left(\int_{B_{R_1}}|u_R-u_{R'}|(x,t)\,dx\right)^{1-m}\le&\left(\int_{B_{2R_1}}|u_R-u_{R'}|(x,0)\,dx\right)^{1-m}+CR_1^{n(1-m)-2}t\\ =&CR_1^{n(1-m)-2}t\quad\forall R>2R_1,R'>2R_1>0,t>0\\ \Rightarrow\qquad\int_{B_{R_1}}|u_R-u|(x,t)\,dx\le&C'R_1^{n-\frac{2}{1-m}}t\quad\forall R>2R_1>0,t>0\quad\text{as } R'\to\infty.\end{aligned}\tag{2.24}$$

Hence

$$\begin{aligned}\int_{B_{R_1}}|u(x,t)-u_0(x)|\,dx\le&\int_{B_{R_1}}|u_R(x,t)-u_0(x)|\,dx+\int_{B_{R_1}}|u-u_R|(x,t)\,dx\\ \le&\int_{B_{R_1}}|u_R(x,t)-u_0(x)|\,dx+C'R_1^{n-\frac{2}{1-m}}t\quad\forall R>2R_1>0,t>0.\end{aligned}\tag{2.25}$$

Letting $t\to 0$ in (2.25),

$$\begin{aligned}\limsup_{t\to 0}\int_{B_{R_1}}|u(x,t)-u_0(x)|\,dx\le&0\quad\forall R_1>0\\ \Rightarrow\qquad\lim_{t\to 0}\int_{B_{R_1}}|u(x,t)-u_0(x)|\,dx=&0\quad\forall R_1>0.\end{aligned}$$

Hence u is the unique solution of (1.5) in $\mathbb{R}^n\times(0,T)$.
Second method: By (2.4), (2.19), (2.21), and (2.23), the sequence $\{w_R:R>R_0\}$ are uniformly bounded below and above on every compact subset of $\mathbb{R}^n\times(0,T)$. Hence the equation (1.1) for $\{w_R:R>R_0\}$ are uniformly parabolic on every compact subset of $\mathbb{R}^n\times(0,T)$. By the parabolic Schauder estimates [LSU] the sequence $\{w_R:R>R_0\}$ are equi-Holder continuous in C^2 on on every compact subset of $\mathbb{R}^n\times(0,T)$. Hence w_R increases and converges uniformly on every compact subset of $\mathbb{R}^n\times(0,T)$ to a solution w of (1.1) as $R\to\infty$. By an argument similar to the first method of proof w has initial value u_0. Hence by the uniqueness of solution $w=u$ is the unique solution of (1.5) in $\mathbb{R}^n\times(0,T)$. □

Theorem 2.6. *Let $n\ge 3$, $0<m\le(n-2)/n$, and let p satisfy* (1.3). *Let $C_1>0$ be as in Lemma 2.5. Suppose $0\le u_0\in L^p_{loc}(\mathbb{R}^n)$ satisfies* (1.10) *for some constant $T>0$. Let $0\le g_R\in L^\infty(\partial B_R\times(0,T))$ and let $\widetilde{u}_R$ be the solution of* (1.13) *in Q_R^T with $g=g_R$ and $\Omega=B_R$. Then $\widetilde{u}_R$ converges uniformly on every compact subset of $\mathbb{R}^n\times(0,T)$ to the unique solution u of* (1.5) *as $R\to\infty$.*

Proof: Let R_0, w_R, be given by Lemma 2.5 and let u_R be as in the proof of Theorem 1.1. By the maximum principle $u_R\ge\widetilde{u}_R\ge w_{R/5}$ in Q_R^T for any $R>5R_0$. Since both u_R and $w_{R/5}$ converges uniformly on every compact subset of $\mathbb{R}^n\times(0,T)$ to the unique solution of (1.5) as $R\to\infty$, $\widetilde{u}_R$ converges uniformly on every compact subset of $\mathbb{R}^n\times(0,T)$ to the unique solution u of (1.5) in $\mathbb{R}^n \times (0, \infty)$ which satisfies (1.11) in $\mathbb{R}^n \times (0, \infty)$ as $R \to \infty$ and the theorem follows. $\square$

By the proof of Theorem 1.1 and Theorem 2.6, Theorem 1.2 follows. In fact we have the following more general result.

Theorem 2.7. *Let $n \geq 3$, $0 < m \leq (n-2)/n$, and let p satisfy* (1.3). *Let $C_1 > 0$ be as in Lemma 2.5. Suppose $0 \leq u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies* (1.6). *Let $0 \leq g_R \in L^\infty(\partial B_R \times (0, \infty))$ and let $\widetilde{u}_R$ be the solution of* (1.13) *in $Q_R^{T_R}$ with $g = g_R$ and $\Omega = B_R$ where T_R is the extinction time of $\widetilde{u}_R$ with $T_R = \infty$ if $\widetilde{u}_R > 0$ in Q_R. Then $\widetilde{u}_R$ converges uniformly on every compact subset of $\mathbb{R}^n \times (0, \infty)$ to the unique solution u of* (1.5) *in $\mathbb{R}^n \times (0, \infty)$ which satisfies* (1.11) *in $\mathbb{R}^n \times (0, \infty)$ as $R \to \infty$.*

Corollary 2.8. *Let $n \geq 3$, $0 < m \leq (n-2)/n$, $q < 2/(1-m)$, and p satisfy* (1.3). *Suppose $0 \leq u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies $u_0 = \overline{u}_0 + \phi$ where $0 \leq \overline{u}_0 \in L^p_{loc}(\mathbb{R}^n)$ and $\phi \in L^1(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$ such that*

$$\liminf_{|x| \to \infty} |x|^q \overline{u}_0(x) \geq A \tag{2.26}$$

for some constant $A > 0$. Then (1.5) *has a unique global solution u in $\mathbb{R}^n \times (0, \infty)$ which satisfies* (1.11) *in $\mathbb{R}^n \times (0, \infty)$.*

Proof: By (2.26) there exists a constant $R_1 > 0$ such that

$$\overline{u}_0(x) \geq (A/2)|x|^{-q} \quad \forall |x| \geq R_1.$$

Since $q < 2/(1-m) \leq n$,

$$\begin{aligned}
\frac{1}{R^{n-\frac{2}{1-m}}} \int_{B_R} u_0 \, dx \geq& \frac{1}{R^{n-\frac{2}{1-m}}} \int_{B_R} \overline{u}_0 \, dx - \frac{1}{R^{n-\frac{2}{1-m}}} \|\phi\|_{L^1} \\
\geq& \frac{\omega_n A}{2(n-q)} (R^{\frac{2}{1-m}-q} - R^{\frac{2}{1-m}-n} R_1^{n-q}) - R^{\frac{2}{1-m}-n} \|\phi\|_{L^1} \quad \forall R > R_1 \\
\to& \infty \quad \text{as } R \to \infty.
\end{aligned}$$

Hence by Theorem 1.2 (1.5) has a unique global solution u in $\mathbb{R}^n \times (0, \infty)$ which satisfies (1.11) in $\mathbb{R}^n \times (0, \infty)$ and the corollary follows. $\square$

Corollary 2.9. *Let $n \geq 3$, $0 < m \leq (n-2)/n$, $q < 2/(1-m)$, and p satisfy* (1.3). *Suppose $0 \leq u_0 \in L^p_{loc}(\mathbb{R}^n)$ satisfies $u_0(x) \geq C_0/|x|^q$ for any $|x| \geq R_1$ and some constants $C_0 > 0$, $R_1 > 0$. Then there exists a unique global solution u of* (1.5) *in $\mathbb{R}^n \times (0, \infty)$ which satisfies* (1.11) *in $\mathbb{R}^n \times (0, \infty)$.*

Lemma 2.10. *Let $n \geq 3$, $0 < m \leq (n-2)/n$, and $q < n/p$ for some constant p satisfying* (1.3). *Let ψ be the unique solution of* (1.5) *in $\mathbb{R}^n \times (0, \infty)$ with initial value $A|x|^{-q}$ which satisfies* (1.11) *in $\mathbb{R}^n \times (0, \infty)$ given by Corollary 2.9. Then*

$$\psi(x, t) \leq A|x|^{-q} \quad \forall x \neq 0, t > 0. \tag{2.27}$$

Proof: For any $R > 0, k > AR^{-q}$, let $v_{R,k}$ be the solution of the problem

$$\begin{cases} v_t = \dfrac{n-1}{m}\Delta v^m & \text{in } Q_R \\ v = AR^{-q} & \text{on } \partial B_R \times (0,\infty) \\ v(x,0) = \min(A|x|^{-q}, k) & \text{in } B_R. \end{cases}$$

Then by the maximum principle (Lemma 2.3 of [DaK]),

$$AR^{-q} \le v_{R,k} \le v_{R,k'} \le k' \quad \text{in } Q_R \quad \forall k' > k > AR^{-q}, R > 0. \tag{2.28}$$

By Theorem 2.3 for any $t_2 > t_1 > 0$, $R > 0$, there exists a constant $M > 0$ such that

$$v_{R,k} \le M \quad \text{in } \overline{B}_R \times [t_1, t_2] \quad \forall k > AR^{-q}. \tag{2.29}$$

By (2.28), (2.29), and an argument similar to the proof of Theorem 1.2, $v_{R,k}$ increases and converges uniformly on every compact subset of $\overline{B}_R \times (0,\infty)$ as $k \to \infty$ to the solution v_R of

$$\begin{cases} v_t = \dfrac{n-1}{m}\Delta v^m & \text{in } Q_R \\ v = AR^{-q} & \text{on } \partial B_R \times (0,\infty) \\ v(x,0) = A|x|^{-q} & \text{in } B_R. \end{cases}$$

Let $t_2 > t_1 > 0$. For any $k_0 > AR^{-q}$ we choose $\varepsilon_0 > 0$ such that $k_0 < A\varepsilon_0^{-q}$. Then by the maximum principle,

$$\begin{aligned} & v_{R,k} \le A|x|^{-q} \quad \forall \varepsilon \le |x| \le R, t > 0, k \ge k_0, 0 < \varepsilon < \varepsilon_0 \\ \Rightarrow \quad & v_R \le A|x|^{-q} \quad \forall 0 < |x| \le R, t > 0 \quad \text{as } k \to \infty. \end{aligned} \tag{2.30}$$

By Theorem 2.7 v_R converges to ψ uniformly on every compact subset of $\mathbb{R}^n \times (0,\infty)$ as $R \to \infty$. Letting $R \to \infty$ in (2.30) we get (2.27) and the lemma follows. □

Theorem 2.11. *Let $n \ge 3$, $0 < m \le (n-2)/n$, and $q < n/p$ for some constant p satisfying* (1.3). *Let α, β, be given by* (1.7). *Then there exists a unique radially symmetric self-similar solution ψ of* (1.5) *in $\mathbb{R}^n \times (0,\infty)$ with initial value $A|x|^{-q}$ which satisfies* (1.11) *and* (2.27) *in $\mathbb{R}^n \times (0,\infty)$ and*

$$\psi(x,t) = \gamma^q \psi(\gamma x, \gamma^{\frac{1}{\beta}} t) \quad \forall x \in \mathbb{R}^n, t > 0, \gamma > 0. \tag{2.31}$$

In particular,

$$\psi(x,t) = t^{-\alpha}\psi(t^{-\beta}x, 1) \quad \forall x \in \mathbb{R}^n, t > 0. \tag{2.32}$$

Moreover $\widetilde{v}(x) = \psi(x,1)$ satisfies (1.9) *in $\mathbb{R}^n$. If in addition $0 < m < (n-2)/n$, then $\psi \in C((\mathbb{R}^n \setminus \{0\}) \times (0,\infty))$ and*

$$|x|^q \psi(x,t) \to A \quad \text{uniformly on } [0,T] \quad \text{as } |x| \to \infty. \tag{2.33}$$

Proof: By Corollary 2.9 and Lemma 2.10 there exists a unique solution ψ of (1.5) in $\mathbb{R}^n\times(0,\infty)$ with initial value $A|x|^{-q}$ which satisfies (1.11) and (2.27) in $\mathbb{R}^n\times(0,\infty)$. Since $\psi(F(x),t)$ is also a solution of (1.1) in $\mathbb{R}^n\times(0,\infty)$ with initial value $A|x|^{-q}$ for any rotation $F:\mathbb{R}^n\to\mathbb{R}^n$, by uniqueness of solution,

$$\psi(x,t)=\psi(F(x),t)\quad\forall x\in\mathbb{R}^n, t\ge 0.$$

Hence $\psi(x,t)$ is radially symmetric in x. For any $\gamma>0$, let $\psi_\gamma(x,t)=\gamma^q\psi(\gamma x,\gamma^{\frac{1}{\beta}}t)$ where β is given by (1.7). Then ψ_γ also satisfies (1.1) in $\mathbb{R}^n\times(0,\infty)$ with initial value $A|x|^{-q}$. By the uniqueness of solution,

$$\psi(x,t)=\psi_\gamma(x,t)\quad\forall x\in\mathbb{R}^n, t>0,\gamma>0$$

and (2.31) follows. Hence ψ is the unique radially symmetric self-similar solution of (1.5) in $\mathbb{R}^n\times(0,\infty)$ with initial value $A|x|^{-q}$ which satisfies (1.11) in $\mathbb{R}^n\times(0,\infty)$. Putting $\gamma=t^{-\beta}$ in (2.31),

$$\psi(x,t)=t^{-\alpha}\psi(t^{-\beta}x,1)=t^{-\alpha}\widetilde{v}(t^{-\beta}x)\quad\forall x\in\mathbb{R}^n, t>0 \tag{2.34}$$

and (2.32) follows. Substituting (2.34) into (1.1), we get that $\widetilde{v}$ satisfies (1.9) in $\mathbb{R}^n$.

We now let $0<m<(n-2)/n$. We will use a modification of the proof of Corollary 2.8 and Corollary 2.9 of [Hu1] to show that ψ satisfies

$$\psi(|x_0|,t)=\psi(x_0,t)\to A|x_0|^{-q}\quad\text{as } t\to 0\quad\forall x_0\ne 0. \tag{2.35}$$

We will first compare ψ with the Barenblatt solution B_k given by (1.4) for some constants $T>0$, $k>0$, to be determined later. Since by (1.3) $q<n/p<2/(1-m)$, by the Young inequality,

$$\begin{aligned}|x|^q=&(T^{\frac{1}{n-2-nm}}|x|)^qT^{-\frac{q}{n-2-nm}}\\ \le&(q(1-m)/2)(T^{\frac{1}{n-2-nm}}|x|)^{\frac{2}{1-m}}+(1-(q(1-m)/2))T^{-\frac{2\alpha}{n-2-nm}}\\ \le&q(1-m)(k+T^{\frac{2}{n-2-nm}}|x|^2)^{\frac{1}{1-m}}\end{aligned} \tag{2.36}$$

where α is given by (1.7) and $k=(2q^{-1}(1-m)^{-1}-1)^{1-m}T^{-\frac{2\alpha(1-m)}{n-2-nm}}$. Let

$$T=\left(\frac{A}{q(1-m)C_*^{\frac{1}{1-m}}}\right)^{\frac{n-2-mn}{n}}$$

where $C_*=2(n-1)(n-2-nm)/(1-m)$. Then by (2.36),

$$B_k(x,0)\le A|x|^{-q}\quad\forall x\in\mathbb{R}^n. \tag{2.37}$$

Let u_R be the solution of (1.15) in Q_R with $\Omega=B_R$ and initial value $A|x|^{-q}$. By (2.37) and the maximum principle ([Hu2]),

$$u_R(x,t)\ge B_k(x,t)\quad\text{in } Q_R. \tag{2.38}$$

Since by the proof of Theorem 1.1 u_R converges uniformly on every compact subset of $\mathbb{R}^n \times (0,\infty)$ to ψ as $R \to \infty$, letting $R \to \infty$ in (2.38),

$$\psi(x,t) \ge B_k(x,t) \quad \forall x \in \mathbb{R}^n, t > 0. \tag{2.39}$$

For any $x_0 \neq 0$, let $R_1 = |x_0|/2$. By (2.27) and (2.39) there exist constants $C_2 > 0$, $C_3 > 0$, such that

$$C_2 \le \psi(x,t) \le C_3 \quad \forall x \in B_{R_1}(x_0), t > 0. \tag{2.40}$$

Since $A|x|^{-q}$ is continuous for all $x \neq 0$, by (2.40) and an argument similar to that of [DaFK] (2.35) follows. Hence putting $\gamma = 1/|x|$, $x \neq 0$, into (2.31) by (2.35) we have

$$|x|^q \psi(x,t) = \psi(1, t/|x|^{\frac{1}{\beta}}) \to A$$

uniformly on $[0,T]$ as $|x| \to \infty$ for any $T > 0$ and the theorem follows. □

3 Asymptotic behaviour of solutions

In this section we will prove the asymptotic large time behaviour of the global solution of (1.5).

Proof of Theorem 1.3: We will use a modification of the proof Theorem 2.1 of [Hs1] to prove the theorem. For any $\gamma \ge 1$, let $u_\gamma(x,t) = \gamma^q u(\gamma x, \gamma^{\frac{1}{\beta}} t)$, $u_{0,\gamma}(x) = \gamma^q u_0(\gamma x)$, $\overline{u}_{0,\gamma}(x) = \gamma^q \overline{u}_0(\gamma x)$, and $\phi_\gamma(x) = \gamma^q \phi(\gamma x)$. Let $C_1 > 0$ be as in the proof of Lemma 2.5 and Theorem 1.1. Then

$$u_\gamma(x,0) = u_{0,\gamma}(x) = \overline{u}_{0,\gamma}(x) + \phi_\gamma(x). \tag{3.1}$$

By the same computation as the proof of Theorem 2.1 of [Hs1] we get

$$\|\phi_\gamma\|_{L^p} \le \|\phi\|_{L^p} \quad \forall \gamma \ge 1, \tag{3.2}$$

$$\int_{B_R} \overline{u}_{0,\gamma}^p \, dx \le C(1 + R^{n-pq}) \quad \forall R > 0, \gamma \ge 1, \tag{3.3}$$

and

$$\|\phi_\gamma\|_{L^1} \le \gamma^{q-n} \|\phi\|_{L^1} \to 0 \quad \text{as } \gamma \to \infty. \tag{3.4}$$

By (1.12) for any $0 < \varepsilon < A$ there exists a constant $R_\varepsilon > 0$ such that

$$\begin{aligned} & (A-\varepsilon)|x|^{-q} \le \overline{u}_0(x) \le (A+\varepsilon)|x|^{-q} \quad \forall |x| \ge R_\varepsilon \\ \Rightarrow \quad & (A-\varepsilon)|x|^{-q} \le \overline{u}_{0,\gamma}(x) \le (A+\varepsilon)|x|^{-q} \quad \forall |x| \ge R_\varepsilon/\gamma. \end{aligned} \tag{3.5}$$

Hence by (3.3) and (3.5),

$$
\begin{aligned}
&\int_{|x|\le R} |\overline{u}_{0,\gamma}(x) - A|x|^{-q}|\,dx \\
\le&\varepsilon A \int_{R_\varepsilon/\gamma\le|x|\le R} |x|^{-q}\,dx + C(R_\varepsilon/\gamma)^{n(1-\frac{1}{p})}\left(\int_{|x|\le R_\varepsilon/\gamma} \overline{u}_{0,\gamma}^{p}\,dx\right)^{\frac{1}{p}} + A\int_{|x|\le R_\varepsilon/\gamma} |x|^{-q}\,dx \\
\le&\varepsilon A \int_{|x|\le R} |x|^{-q}\,dx + C(R_\varepsilon/\gamma)^{n(1-\frac{1}{p})}(1 + (R_\varepsilon/\gamma)^{n-pq})^{\frac{1}{p}} + A\int_{|x|\le R_\varepsilon/\gamma} |x|^{-q}\,dx \quad \forall R > 0, \gamma > R_\varepsilon/R.
\end{aligned}
\tag{3.6}
$$

Letting first $\gamma \to \infty$ and then $\varepsilon \to 0$ in (3.6), by (3.4) we get

$$
\begin{aligned}
&\lim_{\gamma\to\infty} \int_{|x|\le R} |\overline{u}_{0,\gamma}(x) - A|x|^{-q}|\,dx = 0 \quad \forall R > 0 \\
\Rightarrow\quad &\lim_{\gamma\to\infty} \int_{|x|\le R} |u_{0,\gamma}(x) - A|x|^{-q}|\,dx = 0 \quad \forall R > 0.
\end{aligned}
\tag{3.7}
$$

By (3.1), (3.2), (3.3), and Corollary 2.2, for any constants $R > 0$, $t_2 > t_1 > 0$, there exists a constant $C_2 > 0$ such that

$$
u_\gamma(x,t) \le C_2 \quad \forall |x| \le R, t_1 \le t \le t_2. \tag{3.8}
$$

Let δ, R_0, be as in Lemma 2.5. Let $t_2 > t_1 > 0$. Then by (3.4),

$$
\begin{aligned}
\frac{1}{R^{n-\frac{2}{1-m}}}\int_{|x|\le R} u_{0,\gamma}\,dx \ge& \frac{1}{R^{n-\frac{2}{1-m}}}\left(\int_{|x|\le R} A|x|^{-q}\,dx - \int_{|x|\le R} |\overline{u}_{0,\gamma}(x) - A|x|^{-q}|\,dx - \|\phi_\gamma\|_{L^1(\mathbb{R}^n)}\right) \\
\ge& \frac{A\omega_n}{n-q}R^{\frac{2}{1-m}-q} - \frac{1}{R^{n-\frac{2}{1-m}}}\left(\int_{|x|\le R} |\overline{u}_{0,\gamma}(x) - A|x|^{-q}|\,dx + \gamma^{q-n}\|\phi\|_{L^1(\mathbb{R}^n)}\right).
\end{aligned}
\tag{3.9}
$$

Since $q < n/p < 2/(1-m)$, there exists a constant $R_1 > R_0$ such that

$$
\frac{A\omega_n}{n-q}R^{\frac{2}{1-m}-q} \ge \frac{2C_1 t_2^{\frac{1}{1-m}}}{(1-\delta)^{n-\frac{2}{1-m}}} \quad \forall R \ge R_1. \tag{3.10}
$$

Let $R \ge R_1$. By (3.7) there exists $\gamma_R > 1$ such that the last term on the right hand side of (3.9) is less than

$$
\le C_1 t_2^{\frac{1}{1-m}} \quad \forall \gamma \ge \gamma_R. \tag{3.11}
$$

Let $|x_0| \le \delta R$. Then by (3.9), (3.10), and (3.11),

$$
\frac{1}{R^{n-\frac{2}{1-m}}}\int_{B_R(x_0)} u_{0,\gamma}\,dx \ge \frac{1}{R^{n-\frac{2}{1-m}}}\int_{|x|\le(1-\delta)R} u_{0,\gamma}\,dx \ge C_1 t_2^{\frac{1}{1-m}} \quad \forall R \ge \frac{R_1}{1-\delta}, \gamma \ge \gamma_R. \tag{3.12}
$$

Let $v_{R,\gamma}$ be the weak solution of (2.3) in $Q_{5R}^{T_{R,\gamma}}$ with u_0 being replaced by $u_{0,\gamma}$ where $T_{R,\gamma} > 0$ is the extinction time of $v_{R,\gamma}$. Let $\widetilde{v}_{R,\gamma}$ be the weak solution of (2.6) in $Q_{3R}^{\widetilde{T}_{R,\gamma}}(x_0)$ with u_0 being replaced by $u_{0,\gamma}$ where $\widetilde{T}_{R,\gamma} > 0$ is the extinction time of $\widetilde{v}_{R,\gamma}$. By (3.3) and an argument similar to that on P.232-233 of [P] there exists a constant $C > 0$ such that

$$\widetilde{T}_{R,\gamma} \le C\left(\int_{B_{2R}} u_{0,\gamma}^p \, dx\right)^{\frac{2}{n}} \le C'(1 + R^{n-pq})^{\frac{2}{n}} \quad \forall R > 0, \gamma \ge 1. \tag{3.13}$$

Since by the maximum principle $u_\gamma \ge v_{R,\gamma} \ge \widetilde{v}_{R,\gamma}$ in $Q_{3R}^{\widetilde{T}_{R,\gamma}}(x_0)$, by (3.12), (3.13), and an argument similar to the proof of Lemma 2.5,

$$\widetilde{T}_{R,\gamma} > t_2 \quad \forall R \ge R_1/(1-\delta), \gamma \ge \gamma_R,$$

and there exist constants $C_0' > 0$, $C_0'' > 0$, such that

$$\begin{aligned}\inf_{\overline{B_{\delta R}}\times[t_1,t_2]} u_\gamma^m \ge& \inf_{\overline{B_{\delta R}}\times[t_1,t_2]} v_{R,\gamma}^m \\ \ge& C_0' \widetilde{T}_{R,\gamma}^{-\frac{1}{1-m}} R^{-\frac{2m}{1-m}} t_1^{\frac{m}{1-m}} (2t_2^{\frac{m}{1-m}} - t_1^{\frac{m}{1-m}}) \\ \ge& C_0''(1 + R^{n-pq})^{-\frac{2}{n(1-m)}} R^{-\frac{2m}{1-m}} t_1^{\frac{m}{1-m}} (2t_2^{\frac{m}{1-m}} - t_1^{\frac{m}{1-m}}) > 0 \quad \forall R \ge \frac{R_1}{1-\delta}, \gamma \ge \gamma_R.\end{aligned} \tag{3.14}$$

By (3.8) and (3.14) the equation (1.1) for the family of functions $\{u_\gamma : \gamma \ge \gamma_R\}$ are uniformly parabolic on $\overline{B_{\delta R}} \times [t_1, t_2]$ for any $R \ge R_1/(1-\delta)$. By the parabolic Schauder estimates [LSU] the family of functions $\{u_\gamma : \gamma \ge \gamma_R\}$ are equi-Holder continuous on $\overline{B_{\delta R}} \times [t_1, t_2]$ for any $R \ge R_1/(1-\delta)$.

Let $\{\gamma_i\}_{i=1}^\infty$ be a sequence such that $\gamma_i \ge 1$ for all $i \in \mathbb{Z}^+$ and $\gamma_i \to \infty$ as $i \to \infty$. Then by the Ascoli Theorem and a diagonalization argument the sequence $\{u_{\gamma_i}\}_{i=1}^\infty$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges uniformly on every compact subset of $\mathbb{R}^n \times (0,\infty)$ as $i \to \infty$ to some solution ψ of (1.1) that satisfies (1.11) in $\mathbb{R}^n \times (0,\infty)$. We will now prove that ψ has initial value $A|x|^{-q}$. For any $i, j \in \mathbb{Z}^+$, $R > 0$, $t > 0$,

$$\begin{aligned}&\int_{|x|\le R} |\psi(x,t) - A|x|^{-q}| \, dx \\ \le& \int_{|x|\le R} |\psi(x,t) - u_{\gamma_j}(x,t)| \, dx + \int_{|x|\le R} |u_{\gamma_j}(x,t) - u_{0,j}| \, dx + \int_{|x|\le R} |u_{0,j} - A|x|^{-q}| \, dx.\end{aligned} \tag{3.15}$$

Then similar to (2.24) we have

$$\begin{aligned}&\left(\int_{|x|\le R} |u_{\gamma_i} - u_{\gamma_j}|(x,t) \, dx\right)^{1-m} \le \left(\int_{|x|\le 2R} |u_{0,\gamma_i} - u_{0,\gamma_j}| \, dx\right)^{1-m} + CR^{n(1-m)-2} t \\ \Rightarrow \quad& \int_{|x|\le R} |u_{\gamma_i} - u_{\gamma_j}|(x,t) \, dx \le C'\left(\int_{|x|\le 2R} |u_{0,\gamma_i} - u_{0,\gamma_j}| \, dx + R^{n-\frac{2}{1-m}} t^{\frac{1}{1-m}}\right) \quad \forall i, j \in \mathbb{Z}^+, R > 0, t > 0.\end{aligned} \tag{3.16}$$

Letting $i \to \infty$ in (3.16), by (3.7) we get

$$\int_{|x|\le R} |\psi - u_{\gamma_j}|(x,t)\,dx \le C'\left(\int_{|x|\le 2R} |A|x|^{-q} - u_{0,\gamma_j}|\,dx + R^{n-\frac{2}{1-m}} t^{\frac{1}{1-m}}\right) \quad \forall j \in \mathbb{Z}^+, R > 0, t > 0. \tag{3.17}$$

By (3.15) and (3.17),

$$\begin{aligned}&\int_{|x|\le R} |\psi(x,t) - A|x|^{-q}|\,dx\\ \le&\int_{|x|\le R} |u_{\gamma_j}(x,t) - u_{0,j}|\,dx + C\int_{|x|\le 2R} |u_{0,\gamma_j} - A|x|^{-q}|\,dx + CR^{n-\frac{2}{1-m}} t^{\frac{1}{1-m}} \quad \forall j \in \mathbb{Z}^+, R > 0, t > 0.\end{aligned} \tag{3.18}$$

Letting first $t \to 0$ and then $j \to \infty$ in (3.18), by (3.7) we get

$$\lim_{t\to 0}\int_{|x|\le R} |\psi(x,t) - A|x|^{-q}|\,dx \quad \forall R > 0.$$

Hence ψ has initial value $A|x|^{-q}$. By Theorem 2.11 ψ is the unique self-similar radially symmetric solution of (1.1) in $\mathbb{R}^n \times (0,\infty)$ with initial value $A|x|^{-q}$. Since the sequence $\{\gamma_i\}_{i=1}^{\infty}$ is arbitrary, u_γ converges uniformly to ψ as $\gamma \to \infty$ on every compact subset of $\mathbb{R}^n \times (0,\infty)$. In particular

$$u_\gamma(x,1) = \gamma^q u(\gamma x, \gamma^{\frac{1}{\beta}}) \to \psi(x,1) \tag{3.19}$$

uniformly on every compact subset of $\mathbb{R}^n \times (0,\infty)$ as $\gamma \to \infty$. By (3.19),

$$v(x,t) = t^\alpha u(t^\beta x, t) \to \psi(x,1)$$

uniformly on every compact subset of $\mathbb{R}^n \times (0,\infty)$ as $t \to \infty$ and the theorem follows. □

References

[A] D.G. Aronson, *The porous medium equation*, CIME Lectures, in Some problems in Nonlinear Diffusion, Lecture Notes in Mathematics 1224, Springer-Verlag, New York, 1986.

[AC] D.G. Aronson and L.A. Caffarelli, *The initial trace of a solution of the porous medium equation*, Trans. Amer. Math. Soc. 280 (1983), no. 1, 351–366.

[BBDGV] A. Blanchet, M. Bonforte, J. Dolbeault, G. Grillo and J.L. Vazquez, *Asymptotics of the fast diffusion equation via entropy estimates*, Arch. Rat. Mech. Anal. 191 (2009), 347–385.

[BV] M. Bonforte and J.L. Vazquez, *Positivity, local smoothing, and Harnack inequalities for very fast diffusion equations*, Advances in Math. 223 (2010), 529–578.

[BC] P. Bénlian and M.G. Crandall, *Regularizing effects of homogenous evolution equations*, pp. 23–39, in Contributions to Analysis and Geometry (suppl. to Amer. J. Math.), Johns Hopkins Univ. Press, Baltimore, MD, 1981.

[DaFK] B.E.J. Dahlberg, E. Fabes and C.E. Kenig, *A Fatou thoerem for solutions of the porous medium equations*, Proc. Amer. Math. Soc. 91 (1984), 205–212.

[DaK] B.E.J. Dahlberg and C.E. Kenig, *Nonnegative solutions to the generalized porous medium equation*, Rev. Mat. Iberoamericana 2 (1986), 267–305.

[DK] P. Daskalopoulos and C.E. Kenig, *Degenerate diffusion-initial value problems and local regularity theory*, Tracts in Mathematics 1, European Mathematical Society, 2007.

[DP] P. Daskalopoulos and M. Del Pino, *On Nonlinear parabolic equations of very fast diffusion*, Arch. Rat. Mech. Anal. 137 (1997), 363–380.

[DS1] P. Daskalopoulos and N. Sesum, *On the extinction profile of solutions to fast diffusion*, J. Reine Angew Math. 622 (2008), 95–119.

[DS2] P. Daskalopoulos and N. Sesum, *The classification of locally conformally flat Yamabe solitons*, http://arxiv.org/abs/1104.2242.

[HP] M.A. Herrero and M. Pierre, *The Cauchy problem for $u_t = \Delta u^m$ for $0 < m < 1$*, Trans. Amer. Math. Soc. 291 (1985), no. 1, 145–158.

[Hs1] S.Y. Hsu, *Large time behaviour of solutions of a singular diffusion equation in $\mathbb{R}^n$*, Nonlinear Analysis TMA 62 (2005), no. 2, 195–206.

[Hs2] S.Y. Hsu, *Singular limit and exact decay rate of a nonlinear elliptic equation*, http://arxiv.org/abs/1107.2735v1.

[Hu1] K.M. Hui, *On some Dirichlet and Cauchy problems for a singular diffusion equation*, Differential Integral Equations 15 (2002), no. 7, 769V804.

[Hu2] K.M. Hui, *Singular limit of solutions of the very fast diffusion equation*, Nonlinear Anal. TMA 68 (2008), 1120–1147.

[K] T. Kato, *Schrödinger operators with singular potentials*, Israel J. Math. 13 (1973), 135–148.

[LSU] O.A. Ladyzenskaya, V.A. Solonnikov and N.N. Uraltceva, *Linear and quasilinear equations of parabolic type*, Transl. Math. Mono. vol. 23, Amer. Math. Soc., Providence, R.I., U.S.A., 1968.

[P] L.A. Peletier, *The porous medium equation in Applications of Nonlinear Analysis in the Physical Sciences*, H.Amann, N.Bazley, K.Kirchgassner editors, Pitman, Boston, 1981.

[PS] M. Del Pino and M. Sáez, *On the extinction profile for solutions of* $u_t = \Delta u^{(N-2)/(N+2)}$, Indiana Univ. Math. J. 50 (2001), no. 1, 611–628.

[R] G. Rosen, *Nonlinear heat conduction in solid* H_2, Rhys. Rev. B 19 (1979), 2398–2399.

[V1] J.L. Vazquez, *Nonexistence of solutions for nonlinear heat equations of fast-diffusion type*, J. Math. Pures Appl. 71 (1992), 503–526.

[V2] J.L. Vazquez, *Smoothing and decay estimates for nonlinear diffusion equations*, Oxford Lecture Series in Mathematics and its Applications 33, Oxford University Press, Oxford, 2006.

[V3] J.L. Vazquez, *The porous medium equation-Mathematical Theory*, Oxford Mathematical Monographs, Oxford University Press, 2007.